



Hill, R. A. and Sutherland, A. (2017) Hot off the press. Natural Product Reports, (doi:10.1039/c7np90017a).

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

<http://eprints.gla.ac.uk/141464/>

Deposited on: 30 May 2017

Enlighten – Research publications by members of the University of Glasgow
<http://eprints.gla.ac.uk>

Hot off the Press


Robert A. Hill and Andrew Sutherland


School of Chemistry, Glasgow University, Glasgow, UK, G12 8QQ.

E-mail: Bob.Hill@glasgow.ac.uk, Andrew.Sutherland@glasgow.ac.uk


Abstract: A personal selection of 32 recent papers is presented covering various aspects of current developments in bioorganic chemistry and novel natural products such as macrophilone A from *Macrorhynchia philippina*.


A method for validating the structures of natural products by computed ^{13}C NMR chemical shifts of carbons bearing heavy atoms has been developed using a large number of halogenated natural products.¹ This technique, together with computed spin-spin coupling constants, has been used to revise the structure of 16 natural products such as tristichone C that was revised from **1** to **2**. Several iodinated meroterpenoids, including iodocallophycoic acid A **3**, have been isolated from a *Callophycus* sp.² Iodocallophycoic acid A **3**, showed moderate antibiotic activity against MRSA and VREF. A metabolite of a soil-derived *Actinomadura* sp., nonthmicin **4**, is a polyether polyketide with the first example of a natural chlorinated tetronic acid.³






Macrophilone A **5** is a pyrroloiminoquinone from the stinging hydroid *Macrorhynchia philippina*.⁴ The structure of macrophilone A **5** was predicted using long-range NMR correlations and DFT calculations and confirmed by synthesis. Terrazoanthine A **6**, from the zoantharian *Terrazoanthus onoi*, is a representative of a new structural class of 2-aminoimidazole alkaloids.⁵ A biosynthetic pathway to terrazoanthine A **6**, involving a [4+2] cycloaddition, has been proposed. Meyeniin A **7**, from tubers of *Lepidium meyenii* (maca), is a novel sulfur-containing hexahydroimidazo[1,5-*c*]thiazole derivative.⁶ The structure of meyeriin A **7** was confirmed by X-ray analysis and biomimetic synthesis. Macahydantoin A **8**, a thiohydantoin derivative with a new skeleton, was also isolated from *Lepidium meyenii*.⁷ The structure of macahydantoin A **8** was confirmed by synthesis. The C-terminus peptide modification found in leupeptazin **9**, a metabolite of a *Streptomyces* sp., has not been found in nature before.⁸ Leupeptazin **9** shows moderate inhibitory activity towards the collagenase cathepsin K.






A sponge symbiotic association of *Plakortis halichondrioides* and *Xestospongia deweerdtiae* is the source of an inseparable mixture of plakortinic acids A **10** and B **11** that have an unusual bicyclo[4.2.0]octane unit.⁹ It is proposed that this ring system is formed from a linear tetraene precursor. The mixture of plakortinic acids A **10** and B **11** shows cytotoxic activity at submicromolar concentration. The structure of spiroaspertrione A **12**, a metabolite of an *Aspergillus* sp. with a novel terpene-polyketide skeleton, was confirmed by X-ray analysis.¹⁰ A biosynthetic pathway to spiroaspertrione A **12**, involving its co-metabolite andiconin B **13**, has been proposed. Spiroaspertrione A **12** shows potent resensitisation of oxacillin against MRSA.





Fragilolide A **14**, from the gorgonian coral *Junceella fragilis*, has a new 18-norcembrane diterpenoid skeleton with 4,13 and 7,11 cyclisations.¹¹ Norcembranoids are often found in *Sinularia* species of soft corals. This is the first example of a norcembranoid from a gorgonian species. Expression of this gene in *E. coli* produced lydicene **15** that has a new diterpenoid skeleton. A sesterterpene synthase has been

identified from the genome of *Arabidopsis thaliana* that produces the sesterterpenoid thalianatriene **15** which has a novel tricyclic skeleton.¹² A biosynthetic pathway to thalianatriene **15** has been proposed. The structure of bufospirostenin A **16**, from the toad *Bufo bufo gargarizans*, was confirmed by X-ray analysis.¹³ Bufospirostenin A **16** is the first spirostanol to be found in animals.




Fallaxosides B₁ and D₃ are triterpenoid glycosides, from the sea cucumber *Cucumaria fallax*, with the novel aglycones **17** and **18**, respectively.¹⁴ It is proposed that the skeleton of **17** is formed by aldol condensation of a 16,24-diketone precursor and that the skeleton of **18** is formed by a pinacol rearrangement of a 7,8-dihydroxy precursor. The structure of the limonoid triconoid A **19**, from *Trichilia connaroides*, was established by X-ray analysis.¹⁵ The authors propose a biosynthetic pathway to the novel rearranged mexicanolide skeleton of triconoid A **19**. Belamcandanin A **20**, from rhizomes of *Belamcanda chinensis*, is the first example of an adduct of a spiroiridal triterpenoid with an isoflavonoid.¹⁶







Through characterisation of the biosynthetic pathway of the fungal polyketide rubratoxin A **21**, the six redox enzymes involved in the key complexity-generating steps have been revealed.¹⁷ These include α -ketoglutarate- and iron(II)-dependent dioxygenases that hydroxylate sp^3 carbons and, a flavin-dependent dehydrogenase that produces the unsaturated lactone. A β -ketoacyl-acyl carrier protein synthase III-like enzyme, PtmR has been shown to transfer a 6-methylsalicylyl moiety from an iterative type I polyketide synthase to an aminocyclopentitol unit during pactamycin (e.g. pactamycin A **22**) biosynthesis.¹⁸ PtmR was found to accept a wide range of *S*-acyl-*N*-acetylcysteamines as substrates, resulting in the generation of a range of novel pactamycin derivatives.




The final key stages of the biosynthesis of marineosins (e.g. marineosin A **23**) have been established with the identification of a novel bifunctional enzyme MarH, which was shown to catalyse condensation of 4-methoxy-2,2'-bipyrrole-5-carboxaldehyde with 2-undecylpyrrole, as well as hydroxylation of the alkyl side-chain of the resulting prodiginine.¹⁹ Genome mining in bacteria and fungi using the EriG protein, a membrane-bound diterpene cyclase from *Hericium erinaceum* has led to the discovery of seven new diterpene cyclases.²⁰ Expression of the cyclases using an engineered *Escherichia coli* strain allowed characterisation of their products, including the new diterpene lycidene **24**, that possesses an unprecedented skeleton.



A detailed study of NovO, a C-methyltransferase that catalyses the regioselective C8-methylation of the coumarin component of the antibiotic novobiocin **25** in *Streptomyces spheroids* has identified the active site residues responsible for binding the substrate.²¹ One of these residues, His120 was also shown to deprotonate the phenolic 7-hydroxyl, locking the substrate in position for the subsequent methyl transfer step. The products of a putative 2-hydroxyphenylthiazoline biosynthetic gene cluster from the genome of *Streptomyces venezuelae* ATCC 10712 include the known 2-hydroxyphenylthiazolines thiazostatin **26** and watasemycin **27**.²² Gene deletion experiments identified a type B radical-SAM methylase homologue that methylates the thiazoline ring of thiazostatin during the final biosynthetic step to form watasemycin.




The last two steps of the biosynthesis of the broad-spectrum natural product antibiotic fosfomicin **28** have been characterised using a combination of in vitro activity reconstitution and crystallographic analysis of the enzymes involved.²³ The reductase Psf3 was shown to convert 2-oxopropylphosphonate to (*S*)-2-hydroxypropylphosphonate, which is then converted to fosfomicin by a Psf4-catalysed epoxidation. A genome-mining approach has led to a comprehensive understanding of the biosynthesis of lipopeptides in myxobacteria.²⁴ Identification of potential producer strains allowed the discovery and isolation of four novel lipopeptide core types (e.g. **29**).





The biocatalytic total synthesis of the bacterial polycyclic tetramate ikarugamycin **30**, a compound which shows anti-leukemic and anti-inflammatory activity has been achieved using three recombinant enzymes, IkaABC.²⁵ The one-pot, three-enzyme process which uses acetyl-CoA, malonyl-CoA and L-ornithine building blocks allows the construction of 15 carbon-carbon and 2 carbon-nitrogen bonds and sets 8 stereogenic centres, yielding ikarugamycin in 9% overall yield. A biohybrid catalyst



composed of palladium(0) nanoparticles and a crosslinked network of aggregated lipase B from *Candida antarctica* have been used for a one-pot cascade process leading to the resolution of secondary alcohols.²⁶ The first stage of the process involves a palladium-catalysed cycloisomerisation of 4-pentynoic acid that produces a lactone that then acts as an acyl donor in the second stage lipase-catalysed kinetic resolution (Scheme 1).



The first example of a biocatalytic *N*-formylation of amines has been reported.²⁷ The process which uses Novozyme 435 (lipase B from *Candida antarctica*) and ethyl formate was shown to be general and scalable for a wide range of aliphatic and aryl amines (Scheme 2). A one-pot two-step conversion of amines to oximes has been developed using a lipase mediated perhydrolysis of ethyl acetate to generate a peracid, which then performs an oxidation to give the oxime (Scheme 3).²⁸ The optimal method utilised lipase B from *Candida antarctica* and gave the majority of products in excellent yields by a simple extraction protocol.





An optimised, multi-gram scale chemoenzymatic synthesis of *L-threo*-3-benzyloxyspartate **31** has been reported using methylaspartate ammonia lyase (MAL-384A).²⁹ Modifications such as the use of ammonia and high pH led to >98% conversion and 80% isolated yield of **31** (Scheme 4), which is used as a building block for the preparation of probes to study excitatory amino acid transporters. The cloning and overexpression of hydroxynitrile lyases from *Prunus communis* has led to the identification of *PcHNL5* as an excellent biocatalyst for asymmetric hydrocyanation.³⁰ The enzyme was found to convert a range of aromatic aldehydes to valuable chiral cyanohydrin building blocks in excellent yields and optical purities (Scheme 5).



A one-pot two-step asymmetric synthesis of 2-substituted succinate derivatives has been developed using a rhodium-catalysed diazocoupling, followed by an ene-reductase mediated reduction of the resulting *E*-alkene.³¹ Screening a range of ene-

reductases identified OPR1 from *Lycopersicum esculentum* as effective for the reduction of substrates with bulky *tert*-butyl or benzyl esters (Scheme 6). New potential positron emission tomography (PET) tracers have been developed for the in vivo imaging of cannabinoid receptors (CB₁ and CB₂).³² Fluorinated analogues of 11-hydroxy- Δ^8 -tetrahydrocannabinol (e.g. **32**) were synthesised and shown to retain potent pharmacological properties. The authors propose that [¹⁸F]**32** could be used for overall mapping of CB receptors or for selective blocking and competition experiments.



References

- 1 A. G. Kutateladze and D. S. Reddy, *J. Org. Chem.*, 2017, **82**, 3368.
- 2 S. Lavoie, D. Brumley, T. S. Alexander, C. Jasmin, F. A. Carranza, K. Nelson, C. L. Quave and J. Kubanek, *J. Org. Chem.*, 2017, **82**, 4160.
- 3 Y. Igarashi, N. Matsuoka, Y. In, T. Kataura, E. Tashiro, I. Saiki, Y. Sudoh, K. Duangmal and A. Thamchaipenet, *Org. Lett.*, 2017, **19**, 1406.
- 4 K. Zlotkowski, W. M. Hewitt, P. Yan, H. R. Bokesch, M. L. Peach, M. C. Nicklaus, B. R. O'Keefe, J. B. McMahon, K. R. Gustafson and J. S. Schneekloth Jr., *Org. Lett.*, 2017, **19**, 1726.

- 5 P. O. Guillen, K. B. Jaramillo, G. Genta-Jouve, F. Sinniger, J. Rodriguez and O. P. Thomas, *Org. Lett.*, 2017, **19**, 1558.
- 6 M. Zhou, H.-Y. Ma, Z.-H. Liu, G.-Y. Yang, G. Du, Y.-Q. Ye, G.-P. Li and Q.-F. Hu, *J. Agric. Food Chem.*, 2017, **65**, 1887.
- 7 M.-Y. Yu, X.-J. Qin, L.-D. Shao, X.-R. Peng, L. Li, H. Yang and M.-H. Qiu, *Tetrahedron Lett.*, 2017, **58**, 1684.
- 8 N. Kruglyak, D. E. Williams, H. Chen, S. Law, J. Kaleta, I. Villaneuva, J. E. Davies, R. J. Andersen and D. Brömme, *Bioorg. Med. Chem. Lett.*, 2017, **27**, 1397.
- 9 C. Jiménez-Romero, A. D. Rodríguez and S. Nam, *Org. Lett.*, 2017, **19**, 1486.
- 10 Y. He, Z. Hu, W. Sun, Q. Li, X.-N. Li, H. Zhu, J. Huang, J. Liu, J. Wang, Y. Xue and Y. Zhang, *J. Org. Chem.*, 2017, **82**, 3125.
- 11 W. Cheng, M. Ji, X. Li, J. Ren, F. Yin, L. van Ofwegen, S. Yu, X. Chen and W. Lin, *Tetrahedron*, 2017, **73**, 2518.
- 12 J. Shao, Q.-W. Chen, H.-J. Lv, J. He, Z.-F. Liu, Y.-N. Lu, H.-L. Liu, G.-D. Wang and Y. Wang, *Org. Lett.*, 2017, **19**, 1816.
- 13 H.-Y. Tian, L.-J. Ruan, T. Yu, Q.-F. Zheng, N.-H. Chen, R.-B. Wu, X.-Q. Zhang, L. Wang, R.-W. Jiang and W.-C. Ye, *J. Nat. Prod.*, 2017, **80**, 1182.
- 14 A. S. Silchenko, A. I. Kalinovsky, S. A. Avilov, P. S. Dmitrenok, V. I. Kalinin, D. V. Berdyshev, E. A. Chingizova, P. V. Andryjaschenko, K. V. Minin and V. A. Stonik, *Tetrahedron*, 2017, **73**, 2335.
- 15 G.-C. Wang, Y.-Y. Fan, S. L. Shyaula and J.-M. Yue, *Org. Lett.*, 2017, **19**, 2182.
- 16 G. Ni, G.-R. Shi, J.-Y. Li and D.-Q. Yu, *RSC Adv.*, 2017, **7**, 20160.
- 17 J. Bai, D. Yan, T. Zhang, Y. Guo, Y. Liu, Y. Zou, M. Tang, B. Liu, Q. Wu, S. Yu, Y. Tang and Y. Hu, *Angew. Chem. Int. Ed.*, 2017, **56**, 4782.
- 18 M. E. Abugrain, C. J. Brumsted, A. R. Osborn, B. Philmus and T. Mahmud, *ACS Chem. Biol.*, 2017, **12**, 362.
- 19 W. Lu, P. Kancharla and K. A. Reynolds, *Org. Lett.*, 2017, **19**, 1298.
- 20 Y.-l. Yang, S. Zhang, K. Ma, Y. Xu, Q. Tao, Y. Chen, J. Chen, S. Guo, J. Ren, W. Wang, Y. Tao, W.-B. Yin and H. Liu, *Angew. Chem. Int. Ed.*, 2017, **56**, 4749.

- 21 J. C. Sadler, C.-w. H. Chung, J. E. Mosley, G. A. Burley and L. D. Humphreys, *ACS Chem. Biol.*, 2017, **12**, 374.
- 22 Y. Inahashi, S. Zhou, M. J. Bibb, L. Song, M. M. Al-Bassam, M. J. Bibb and G. L. Challis, *Chem. Sci.*, 2017, **8**, 2823.
- 23 P. Olivares, E. C. Ulrich, J. R. Chekan, W. A. van der Donk and S. K. Nair, *ACS Chem. Biol.*, 2017, **12**, 456.
- 24 C. Burgard, N. Zaburannyi, S. Nadmid, J. Maier, H. Jenke-Kodama, E. Luxenburger, H. S. Bernauer and S. C. Wenzel, *ACS Chem. Biol.*, 2017, **12**, 779.
- 25 C. Greunke, A. Glöckle, J. Antosch and T. A. M. Gulder, *Angew. Chem. Int. Ed.*, 2017, **56**, 4351.
- 26 . T. Görbe, K. P. J. Gustafson, O. Verho, G. Kervefors, H. Zheng, X. Zou, E. V. Johnston and J.-E. Bäckvall, *ACS Catal.*, 2017, **7**, 1601.
- 27 R. E. Patre, S. Mal, P. R. Nilkanth, S. K. Ghorai, S. H. Deshpande, M. El Qacemi, T. Smejkal, S. Pal and B. N. Manjunath, *Chem. Commun.*, 2017, **53**, 2382.
- 28 D. Méndez-Sánchez, I. Lavandera, V. Gotor and V. Gotor-Fernández, *Org. Biomol. Chem.*, 2017, **15**, 3196.
- 29 H. Fu, S. H. H. Younes, M. Saifuddin, P. G. Tepper, J. Zhang, E. Keller, A. Heeres, W. Szymanski and G. J. Poelarends, *Org. Biomol. Chem.*, 2017, **15**, 2341.
- 30 Y.-C. Zheng, J.-H. Xu, H. Wang, G.-Q. Lin, R. Hong and H.-L. Yu, *Adv. Synth. Catal.*, 2017, **359**, 1185.
- 31 Y. Wang, M. J. Bartlett, C. A. Denard, J. F. Hartwig and H. Zhao, *ACS Catal.*, 2017, **7**, 2548.
- 32 C. Zanato, A. Pelagalli, K. F. M. Marwick, M. Piras, S. Dall'Angelo, A. Spinaci, R. G. Pertwee, D. J. A. Wyllie, G. E. Hardingham and M. Zanda, *Org. Biomol. Chem.*, 2017, **15**, 2086.